

PTI Observations of 89 Her

Mark Swain[†], Gerard Van Belle[†], Rachel Akeson[‡]

[†] JPL/Caltech, 4800 Oak Grove Dr., Pasadena CA 91109

[‡] IPAC/Caltech, MS 100-22, Pasadena CA 91125

Abstract

We report on observations of the star 89 Her using the Palomar Testbed Interferometer (PTI). Our observations in the H and K band suggest the presence of a circumstellar dust shell. We discuss the observations in the context of simple models for 89 Her.

1 Introduction

The star 89 Her is an extensively studied post-AGB supergiant with a large infrared excess and a radial velocity signature indicative of a low mass companion. As this star is thought to be undergoing mass loss and is sufficiently bright, it was selected as a candidate for PTI observations with the hope of detecting the warmer component of the circumstellar material. The PTI observations do show very unusual behavior of the V^2 estimator; they imply the diameter of the object increases substantially (~ 2) between the H and K bands. Although there are some differences in the published values for parameters of the system, we have used the values below for the purpose of interpreting our interferometer data.

2 Background

- F2 Ibe - post AGB (due to dust shell)
- Distance = 600 pc
- Luminosity = 3300 L_{solar}
- Radius (primary) = 45 R_{solar}
- Temperature = 6500 K
- Mdot = 1e-8 M_{solar}/yr
- Large IR excess suggests dust shell.
- Negligible extension suggests disk geometry for dust.
- 89 Her apparent diameter is 0.7 mas; predicted $V^2 =$ at 2.2 μm .




Figure 1: Photometry for 89Her showing the IR excess. Separate short and long wavelength fits to the data come up with a stellar temperature of about 5850 K and a dust temperature of about 1100 K.

3 The Observations

Day #	scans	start	stop	V ²	jitter (rads)	WL/spec DN	avg lock brakes
99126	15	9:10	9:57	0.044/0.053	0.507	617.3/37.9	0.53
99144	10	9:35	10:16	0.047/0.055	0.958	568.3/34.5	2.1
99177	9	7:31	8:33	0.050/0.053	1.641	1333./105.	4.22
99178	5	7:32	9:04	0.026/0.059	1.782	1461./77.3	8.2
99239	20	3:08	4:56	0.046/0.054	0.794	1385./57.0	0.3
100123	2	11:29	11:31	0.043/0.046	0.665	1880./73.5	18
100130	1	11:24	11:24	0.037/0.037	0.48	902.1/42.1	17
100140	32	9:48	12:14	0.148/0.296	0.942	448.4/9.34	17.5
100154	5	10:42	10:49	0.049/0.060	0.732	1470./38.2	10.4
100159	12	6:51	7:20	0.022/0.425	0.84	419.2/5.70	14.5
100164	5	9:12	10:32	0.039/0.046	0.936	1617./67.2	12.6
101128	20	10:21	12:07	0.198/0.243	1.01	795.6/19.1	15

Table 1: Summary of PTI observations of 89 Her.




Figure 2: Visibility data as a function of wavelength (microns) for the H and K bands.




Figure 3: Uniform disk diameter inferred from the H and K band data as a function of baseline length.




Figure 4: Uniform disk diameter inferred from the H band data as a function of baseline position angle.

4 Can a Binary Produce the Observed V^2 ?

- Binary orbital solution reported for this system (Walters).
- Probable secondary mass given as 0.09 to 0.07 Msolar.
- If primary and secondary are equal (0.5 Msolar), $R/a = 0.44$.
- Intensity dynamic range of PTI implies $L_{\text{secondary}} \geq L_{\text{primary}}/30$ for the binary V^2 to be detected.
- Thus only similar mass and evolutionary stage binary companion could be detected by PTI.
- Binary test case is $M_{\text{primary}} = M_{\text{secondary}}$, $L_{\text{primary}} = L_{\text{secondary}}$, separation is 2.1 mas.




Figure 5: An estimate of the maximum chromatic V^2 effect due to a possible binary companion to 89 Her. For this we have assumed the most extreme possible case for a binary companion; a secondary of identical mass and luminosity. The result produces a chromatic V^2 substantially different than our observations. This suggests our observations are caused by another effect.

5 Signature of the Dust Disk

- Assuming $T \sim 1/r^2$ from a 6500 K stellar surface until the onset of dust formation, ~ 1000 K dust forms at a radius of about $2.6 R_{\text{star}}$ (0.9 milliarcseconds).
- The combined V^2 of star plus dust is:

$$V_{\text{total}}^2 = \frac{F_{\text{star}} V_{\text{star}}^2 + F_{\text{dust}} V_{\text{dust}}^2}{(F_{\text{star}} + F_{\text{dust}})^2}$$

- Solving for V_{dust}^2 we find corrections to the observed V^2 are small due to the simultaneous constraints of the ratio of dust emission to star emission, baseline evolution of the H band data, and size scale of the dust emission.




Figure 6: An ellipse fitted to the H band data. This suggests an disk rotation axis is inclined to the line of sight of about 25 degrees. The total V^2 have been used in this plot. Corrections for the stellar component could change the size of the disk but not the ellipticity.

V_{total}^2	r=.1	r=.2	r=.4	r=.6	r=.8	r=1.0	r=2.0	r=4	r=6	r=8	r=10
0.05	-9.40	-4.64	-2.26	-1.45	-1.05	-0.80	-0.28	0.06	00.24	0.38	0.51
0.10	-8.79	-4.28	-2.01	-1.24	-0.85	-0.60	-0.05	0.38	0.65	0.89	1.11
0.15	-8.19	-3.92	-1.77	-1.03	-0.64	-0.40	0.18	0.69	1.06	1.39	1.72
0.20	-7.58	-3.56	-1.52	-0.81	-0.44	-0.20	0.40	1.00	1.47	1.90	2.32
0.25	-6.98	-3.20	-1.28	-0.60	-0.24	0.00	0.63	1.31	1.88	2.41	2.93
0.30	-6.37	-2.84	-1.03	-0.39	-0.04	0.20	0.85	1.63	2.28	2.91	3.53
0.35	-5.77	-2.48	-0.79	-0.17	0.17	0.40	1.08	1.94	2.69	3.42	4.14
0.40	-5.16	-2.12	-0.54	0.04	0.37	0.60	1.30	2.25	3.10	3.93	4.74
0.45	-4.56	-1.76	-0.30	0.25	0.57	0.80	1.53	2.56	3.51	4.43	5.35
0.50	-3.95	-1.40	-0.05	0.47	0.78	1.00	1.75	2.88	3.92	4.94	5.95
0.55	-3.34	-1.04	0.20	0.68	0.98	1.20	1.98	3.19	4.33	5.44	6.56

Table 2: Permitted values of dust V^2 show in bold for a range of flux ratios between the disk and the star ($r=F_{\text{disk}}/F_{\text{star}}$) and total V^2 .

6 Conclusions

- We have directly detected the warm dust component around 89 Her.
- There is an indication that the dust distribution is not rotationally symmetric.
- We infer a disk orientation of approximately 25 degrees out of the plane of the sky; better $u-v$ coverage would significantly improve this constraint.
- If the disk and binary system share an angular momentum axis, then the secondary would be well within the accepted radius of the primary.
- The secondary, if it exists, probably plays an important roll in suppling material to the disk.

Acknowledgments

M.S. wishes to thank Len Hillenbrand for suggesting this source and Andy Boden for helpful discussions. The research described in this paper was carried out by then Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration.

References

- [] Allard, F., Hauschildt, P.H., Baraffe, I., Chabrier, G., 1996, *ApJ*, 465, 123
- [] Shenton, M., Evans, A., Williams, P.M., *MNRAS*, 273, 906
- [] Millan-Gabet, R., Schloerb, P.F., Traub, W.A., 2001, *ApJ*, 546, 358
- [] van der Veen, W.E.C.J., Waters, L.B.F.M., Trams, N.R., Matthews, H.E., 1994, *AA*, 285, 551
- [] Waters, L.B.F.M, Waelkens, C., Mayor, M., Trams, N.R., 1993, *AA*, 269, 242